

Referenties

Vastgesteld op: 11-08-2010

Methode: -

Regiehouder: IKNL

Gegenereerd op: 24-06-2024

Bron: <https://palliaweb.nl/richtlijnen-palliatieve-zorg/archief/stervensfase-versie-2022/samenvatting-reutelen>

Inhoudsopgave

REFERENTIES	3
-------------	---

Referenties

Vastgesteld: 11-08-2010 Regiehouder: IKNL

Abarshi E (2009)

Abarshi E, Onwuteaka-Philipsen B, Donker G et al. General practitioner awareness of preferred place of death and correlates of dying in a preferred place: a nationwide mortality follow-back study in The Netherlands. *Journal of Pain and Symptom Management* 2009; 38: 568-577.

Back IN (2001)

Back IN, Jenkins K, Blower A et al. A study comparing hyoscine hydrobromide and glycopyrrolate in the treatment of death rattle. *Palliative Medicine* 2001; 15: 329-336.

Bennett M (2002)

Bennett M, Lucas V, Brennan M et al. Using anti-muscarinic drugs in the management of death rattle: evidence-based guidelines for palliative care. *Palliative Medicine* 2002; 16: 369-374.

Berg van den MR (2002)

Berg van den MR, Berg van den M. Voor de laatste tijd – samen werken aan een goede dood. Kampen: Kok, 2002.

Brandt HE (2006)

Brandt HE, Ooms ME, Deliens L et al. The last two days of life of nursing home patients – a nationwide study on causes of death and burdensome symptoms in the Netherlands. *Palliative Medicine* 2006; 20: 533-540.

Breitbart W (2008)

Breitbart W, Alici Y. Agitation and delirium at the end of life: "We couldn't manage him". *JAMA* 2008; 300: 2898-2910.

Broersen SF (2010)

Broersen SF. Pacemaker vertraagt sterven. *Medisch Contact* 2010; 5: 206-207.

Christ GH (2006)

Christ GH, Christ AE. Current approaches to helping children cope with a parent's terminal illness. *CA: A Cancer Journal for Clinicians* 2006; 56: 197-212.

Clark K (2009)

Clark K, Butler M. Noisy respiratory secretions at the end of life. *Current Opinion in Supportive and Palliative Care* 2009; 3: 120-124.

Clukey L (2008)

Clukey L. Anticipatory mourning: processes of expected loss in palliative care. *International Journal of Palliative Nursing* 2008; 14: 316, 318-25.

Dendaas NR (2002)

Dendaas NR. Prognostication in advanced cancer: nurses' perceptions of the dying process. *Oncology Nursing Forum* 2002; 29: 493-499.

Dom H (1999)

Dom H. Spiritual care, need and pain – recognition and response. *European Journal of Palliative Care* 1999; 6: 87-90.

Donnelly SM (2009)

Donnelly SM, Donnelly CN. The experience of the moment of death in a specialist palliative care unit (SPCU). *Irish Medical Journal* 2009; 102: 143-146.

Dumont I (2008)

Dumont I, Dumont S, Mongeau S. End-of-life care and the grieving process: family caregivers who have experienced the loss of a terminal-phase cancer patient. *Qualitative Health Research* 2008; 18: 1049-1061.

Ellershaw J (2001)

Ellershaw J, Smith C, Overill S et al. Care of the dying: setting standards for symptom control in the last 48 hours of life. *Journal of Pain and Symptom Management* 2001; 21: 12-17.

Ellershaw J (2003)

Ellershaw J, Ward C. Care of the dying patient: the last hours or days of life. *BMJ* 2003; 326: 30-34.

Fallowfield LJ (2002)

Fallowfield LJ, Jenkins VA, Beveridge HA. Truth may hurt but deceit hurts more: communication in palliative care. *Palliative Medicine* 2002; 16: 297-303.

Field D (2007)

Field D, Payne S, Relf Met al. Some issues in the provision of adult bereavement support by UK hospices. *Social Science & Medicine* 2007; 64: 428-438.

Flaherty JH (2004)

Flaherty JH. Urinary incontinence and the terminally ill older person. *Clinical Geriatric Medicine* 2004; 20: 467-475.

Georges JJ (2005)

Georges JJ, Onwuteaka-Philipsen BD, Heide van der A et al. Symptoms, treatment and "dying peacefully" in terminally ill cancer patients: a prospective study. *Supportive Care in Cancer* 2005; 13: 160-68.

Gomes B (1974)

Gomes B, Higginson IJ. Where people die (1974–2030): Past trends, future projections and implications for care. *Palliative Medicine* 2008; 22: 33-41.

Grassi L (2007)

Grassi L. Bereavement in families with relatives dying of cancer. *Current Opinion in Supportive and Palliative Care* 2007; 1: 43-49.

Hanratty JF (1994)

Hanratty JF, Higginson I. *Palliative care in terminal illness*. 2nd Edition, Oxford-New York: Radcliffe Medical Press, 1994.

Holley CK (2009)

Holley CK, Mast BT. The impact of anticipatory grief on caregiver burden in dementia caregivers. *Gerontologist* 2009; 49: 388-396.

Hudson PL (2006)

Hudson PL. How well do family caregivers cope after caring for a relative with advanced disease and how can health professionals enhance their support? *Journal of Palliative Medicine* 2006; 9: 694-703.

Hughes A (2000)

Hughes A, Wilcock A, Corcoran R et al. Audit of three antimuscarinic drugs for managing retained secretions. *Palliative Medicine* 2000; 14: 221-222.

Jackson KC (2004)

Jackson KC, Lipman AG. Drug therapy for delirium in terminally ill patients. *Cochrane Database of Systematic Reviews* 2004; 2: CD004770.

Kaasalainen S (2007)

Kaasalainen S, Brazil K, Ploeg J et al. Nurses' perceptions around providing palliative care for long-term care residents with dementia. *Journal of Palliative Care* 2007; 23: 173-180.

Kåss RM (2003)

Kåss RM, Ellershaw J. Respiratory tract secretions in the dying patient: a retrospective study. *Palliative Medicine* 2003; 26: 897-902.

Keely MP (2004)

Keely MP. Final conversations: survivors' memorable messages concerning religious faith and spirituality. *Health Communication* 2004; 16: 87-104.

Kehl KA (2004)

Kehl KA. Treatment of terminal restlessness: a review of the evidence. *Journal of Pain & Palliative Care Pharmacotherapy* 2004; 18: 5-30.

Kelley AS (2009)

Kelley AS, Reid MC, Miller DH et al. Implantable cardioverter-defibrillator deactivation at the end of life: a physician survey. *American Heart Journal* 2009; 157: 702-708.

Klinkenberg M (2004)

Klinkenberg M, Willems DL, Wal van der G et al. Symptom burden in the last week of life. *Journal of Pain and Symptom Management* 2004; 27: 5-13.

Kwan C (2002)

Kwan C. Families' experiences of the last office of deceased family members in the hospice setting. *International Journal of Palliative Nursing* 2002; 8: 266-275.

Legrand M (1998)

Legrand M, Gomas JM. Death rituals: Being present at the last moments of life. *European Journal of Palliative Care* 1998; 5: 191-194.

Lichtenthal WG (2009)

Lichtenthal WG, Nilsson M, Zhang B et al. Do rates of mental disorders and existential distress among advanced stage cancer patients increase as death approaches? *Psycho-Oncology* 2009; 18: 50-61.

Liossi C (1997)

Liossi C, Hatira P, Mystakidou K. The use of the genogram in the palliative care. *Palliative Medicine* 1997; 11: 455-461.

Lunney JR (2003)

Lunney JR, Lynn J, Foley DJ et al. Patterns of functional decline at the end of life. *JAMA* 2003; 289: 2387-2392.

Maguire P (2002)

Maguire P, Pitceathly C. Key communication skills and how to acquire them. *BMJ* 2002; 325: 697-700.

Miguel de Sánchez C (2006)

Miguel de Sánchez C, Elustondo SG, Estirado A et al. Palliative performance status, heart rate and respiratory rate as predictive factors of survival time in terminally ill cancer patients. *Journal of Pain and Symptom Management* 2006; 31: 485-492.

Mitchell G (2008)

Mitchell G. *Palliative Care, a patient-centered approach*. Oxford/New York: Radcliffe Publishing, 2008.

Morita T (1998)

Morita T, Ichiki T, Tsunoda J et al. A prospective study on the dying process in terminally ill cancer patients. *American Journal of Hospice & Palliative Care* 1998; 15: 217-222.

Morita T (2004)

Morita T, Hyodo I, Yoshimi T et al. Incidence and underlying etiologies of bronchial secretion in terminally ill cancer patients: a multicenter, prospective, observational study. *Journal of Pain and Symptom Management* 2004; 27: 533-539.

Parkes CM (1998)

Parkes CM. The dying adult. *BMJ* 1998; 316: 131-1315.

Payne S (2002)

Payne S. Information needs of patients and families. *European Journal of Palliative Care* 2002; 9: 112-114.

Plonk WM Jr (2005)

Plonk WM Jr, Arnold RM. Terminal care: the last weeks of life. *Journal of Palliative Medicine* 2005; 8: 1042-1054.

Roussau P (2007)

Roussau P. Management of symptoms in the actively dying patient. In: Berger AM, Shuster JL, Roenn von JH. *Principles and practice of palliative care and supportive oncology*. 3rd Edition, Ch 57, p 623-632. Philadelphia: Lippincott, Williams & Wilkins, 2007.

Schaap E (2005)

Schaap E, Zuylen van L, Janssen P et al. Reutelen in de stervensfase: een literatuuroverzicht. *Oncologica* 2005; 1: 4-9.

Simon ST (2010)

Simon ST, Higginson IJ, Booth S et al. Benzodiazepines for the relief of breathlessness in advanced malignant and non-malignant diseases in adults. *Cochrane Database of Systematic Review* 2010; 1: CD007354.

Singer PA (1999)

Singer PA, Martin DK, Kelner M. Quality end-of-life care; patients' perspectives. *JAMA* 1999; 281: 163-168.

Shah M (2008)

Shah M, Quill T, Norton S et al. "What bothers you the most?" Initial responses from patients receiving palliative care consultation. *American Journal of Hospice and Palliative Care* 2008; 25: 88-92.

Steinhauser KE (2000)

Steinhauser KE, Clipp EC, McNeilly Met al. In search of a good death: observations of patients, families, and providers. *Annals of Internal Medicine* 2000; 132: 825-832.

Swart SJ (2003)

Swart SJ, Zuylen van L, Rijt van der CC et al. Sterven kost tijd. *Medisch Contact* 2003; 22 : 910-911.

Vachon M (1997)

Vachon M. Burnout: Recent research into staff stress in palliative care. *European Journal of Palliative Care* 1997; 4: 99-103.

Velden van der LFJ (2009)

Velden van der LFJ, Francke AL, Hingstman L et al. Dying from cancer or other chronic diseases in the Netherlands: ten-year trends derived from death certificate data. *BMC Palliative Care* 2009; 8: 4.

Veerbeek L (2007)

Veerbeek L, Zuylen van L, Swart SJ et al. The last 3 days of life in three different care settings in the Netherlands. *Supportive Care in Cancer* 2007; 15: 1117-1123.

Veerbeek L (2008)

Veerbeek L, Zuylen van L, Swart SJ et al. The effect of the Liverpool Care Pathway for the dying: a multi-centre study. *Palliative Medicine* 2008; 22: 145-151.

Veerbeek L (2008)

Veerbeek L, Heide van der A, Vogel-Voogt de E et al. Using the LCP: bereaved relatives' assessments of communication and bereavement. *American Journal of Hospice and Palliative Care* 2008; 25: 207-214.

Wee B (2008)

Wee B, Hillier R. Interventions for noisy breathing in patients near to death. *Cochrane Database of Systematic Reviews* 2008; 1: CD005177.

Wee BL (2006)

Wee BL, Coleman PG, Hillier R et al. The sound of death rattle I: are relatives distressed by hearing this sound? *Palliative Medicine* 2006; 20: 171-175.

Wee BL (2006)

Wee BL, Coleman PG, Hillier R et al. The sound of death rattle II: how do relatives interpret the sound? *Palliative Medicine* 2006; 20: 177-181.

Wildiers H (2002)

Wildiers H, Menten J. Death rattle: prevalence, prevention and treatment. *Journal of Pain and Symptom Management* 2002; 23: 310-317.

Wildiers H (2009)

Wildiers H, Dhaenekint C, Demeulenaere P et al. Atropine, hyoscine butylbromide, or scopolamine are equally effective for the treatment of death rattle in terminal care. *Journal of Pain and Symptom Management* 2009; 38: 124-133.

Zuylen van L (2008)

Zuylen van L, Vos PJ, Veerbeek L et al. Een goed einde: zorgpad verbetert zorg en kwaliteit van leven in de stervensfase. *Medisch Contact* 2008; 503: 2098-2101.